arXiv:1407.1961v3 [nlin.SI] 1 Aug 2015

Singular dynamics of a q-difference Painlevé equation in its initial-value space

N. Joshi and S.B. Lobb

Abstract. We construct the initial-value space of a q-discrete first Painlevé equation explicitly and describe the behaviours of its solutions $w(n)$ in this space as $n \to \infty$, with particular attention paid to neighbourhoods of exceptional lines and irreducible components of the anti-canonical divisor. These results show that trajectories starting in domains bounded away from the origin in initial value space are repelled away from such singular lines. However, the dynamical behaviours in neighbourhoods containing the origin are complicated by the merger of two simple base points at the origin in the limit. We show that these lead to a saddle-point-type behaviour in a punctured neighbourhood of the origin.

Dedicated to Rodney J. Baxter on the occasion of his 75th birthday.

1. Introduction

The discrete Painlevé equations are known to have a simple, beautiful geometric structure [**19**], but very little is known about their general solutions. Motivated by Boutroux's famous study [**5**] of the first Painlevé equation $\mathrm{P_I}$: $y_{tt} = 6\,y^2 - t$, as $|t| \to \infty$, we consider the limit $|\xi| \to \infty$ of its q-discrete version

$$q\mathrm{P_I} : w\big(q\,\xi\big)\,w\big(\xi/q\big) = \frac{w(\xi) - 1/\xi}{w(\xi)^2}, \tag{1.1}$$

where we assume $|q| > 1$ for simplicity. Such equations arise as reductions of Yang-Baxter maps [**2, 18**] and multi-dimensionally consistent integrable lattice equations [**1, 14**].

Starting at a given point in $\mathbb{C}$, repeated iteration of the equation on complex spirals (for $q \in \mathbb{C}$) leads to a sequence of solution values of a discrete Painlevé equation, but no information comparable to that found by Boutroux about global behaviours of solutions of $\mathrm{P_I}$ is available in the plane of the independent variable $\xi \in \mathbb{C}$. Instead, we consider solution trajectories in the space of initial values, which is compactified by embedding in $\mathbb{P}^1 \times \mathbb{P}^1$ and regularized by resolving all singularities[1]. We denote the resulting initial value space by $\mathcal{S}$.

2010 *Mathematics Subject Classification.* 34M30;39A13;34M55.

[1]Note that $\mathbb{P}^1 \times \mathbb{P}^1$ is the two-complex-dimensional space parametrized by pairs of homogeneous coordinates $([x_1 : x_2], [y_1 : y_2])$= $([\lambda\, x_1 : \lambda\, x_2], [\mu\, y_1 : \mu\, y_2])$ for all non-zero $(x_j, y_j) \in \mathbb{C}^2$, $j = 1, 2$, and non-zero λ, $\mu \in \mathbb{C}$.

More precisely, writing $\xi = q^n \xi_0$, $w_n = w(\xi)$, $w_{n+1} = w(q\xi)$, $w_{n-1} = w(\xi/q)$, we study the solutions in $\mathcal{S} = \cup_{n_0 \in \mathbb{C}} S_{n_0}$, where the fibre S_{n_0} is locally described by affine coordinates (w_{n_0}, w_{n_0-1}) and resolved by blowing up 8 base points. Equation (1.1) is symmetric under $w_{n+1} \leftrightarrow w_{n-1}$. In the remainder of the paper, we study the equivalent forward and backward systems

$$\begin{pmatrix} \overline{u} \\ \overline{v} \end{pmatrix} = \begin{pmatrix} \dfrac{q\overline{v} - t}{qu\overline{v}^2} \\ \dfrac{u - t}{u^2 v} \end{pmatrix}, \qquad \begin{pmatrix} \underline{u} \\ \underline{v} \end{pmatrix} = \begin{pmatrix} \dfrac{v - qt}{uv^2} \\ \dfrac{\underline{u} - q^2 t}{\underline{u}^2 v} \end{pmatrix} \tag{1.2}$$

where $t := 1/\xi$, $u_n := w_{2n}$, $v_n := w_{2n-1}$. (Note that bar now denotes $t \mapsto t/q^2$.)

To our knowledge, the term "discrete Painlevé equation" was proposed by Fokas, Its and Kitaev [**9**] for another discrete version of $\mathrm{P_I}$, which arises from transformations of the classical fourth Painlevé equation. The latter equation is an additive-type discrete equation, where coefficient functions are linear in n instead of exponential, i.e., $\xi = \xi_0\, q^n$, as in Equation (1.1).

Many discrete Painlevé equations have now been identified. Following Okamoto's construction for the classical Painlevé equations [**17**], Sakai [**19**] showed how to construct initial value spaces of discrete Painlevé equations as rational surfaces obtained by 9-point resolution of $\mathbb{P}^2$ or equivalently, 8-point resolution of $\mathbb{P}^1 \times \mathbb{P}^1$. In particular, Sakai showed that the initial value space associated with Equation (1.2) is described by the affine Weyl group $A_7^{(1)}$.

Qualitative information about solutions of (1.2) is valuable because Nishioka [**15**] showed that they are higher transcendental functions, which cannot be expressed in terms of basic or q-special functions or in terms of solutions of first-order or linear second-order difference equations. We describe solutions by studying the limit $|\xi| \to \infty$, or equivalently $|t| \to 0$ on its initial value surface.

1.1. Autonomous Limit. In the limit $|t| \to 0$, the system (1.2) becomes

$$\begin{pmatrix} \overline{u} \\ \overline{v} \end{pmatrix} = \begin{pmatrix} v \\ \dfrac{1}{uv} \end{pmatrix}, \qquad \begin{pmatrix} \underline{u} \\ \underline{v} \end{pmatrix} = \begin{pmatrix} \dfrac{1}{uv} \\ u \end{pmatrix} \tag{1.3}$$

or, equivalently $\overline{w}\, w\, \underline{w} = 1$, for either $w = u$ or v. This map is one of the periodic cases of the QRT mappings studied by Tsuda [**20**], which is periodic with period 3, for any initial value. The autonomous system has an invariant given by

$$K(u, v) = \frac{u^2 v^2 + u + v}{uv} \tag{1.4}$$

i.e., $K(\overline{u}, \overline{v}) - K(u, v) = 0 = K(u, v) - K(\underline{u}, \underline{v})$ when u and v satisfy Equations (1.3). The singularities of the invariant curve (i.e., points where the gradient of $K(u, v)$ vanishes) are given by $(u, v) = (\omega, \omega)$ where $\omega^3 = 1$, and $(u, v) = (0, \infty)$, $(u, v) = (\infty, 0)$. These lie on the invariant curve only if $K = 3\omega^2$. Therefore, there are 3 singular fibres in the pencil of curves (1.4).

It is interesting to note that the so called *tronquée* solutions identified by Boutroux for the first Painlevé equation also approach double zeroes of the latter's autonomous invariant. In this sense, the solutions that approach the equilibrium values $(u, v) = (\omega, \omega)$ of the autonomous $q\mathrm{P_I}$ act as discrete analogues of the tronquée solutions. In this paper, however, we focus on the generic solutions of the dynamical system (1.2).

1.2. Fixed points. Fixed points are important for studying limiting solutions. The cases where the solutions of Equation (1.1) approach a steady state as $n \to +\infty$ have been considered in [**12, 16**]. Joshi [**12**] showed that there exists a true solution satisfying $|w| \to 0$ as $n \to +\infty$, which is asymptotic to a divergent asymptotic series expansion in powers of $1/\xi$, but is not a singularity of the limiting invariant K. This vanishing, unstable solution was called the *quicksilver* solution in [**12**] and its trajectory lies in a neighbourhood of the origin in $\mathcal{S}$, which is punctured at two base points (called b_1, b_2 below) that approach the origin. (See Figure 3.2 for a typical such neighbourhood.) Our results show that the dynamics of Equations (1.2) in this punctured neighbourhood resemble that near a saddle point in the initial value space $\mathcal{S}$.

1.3. Outline of the Paper. The structure of the paper is as follows. In Section 2 we construct the initial-value space $\mathcal{S}$ and provide explicit coordinate charts to describe it completely; key points are illustrated here, while full details are given in Appendix A. We also describe the irreducible components of its anti-canonical divisor, with explicit details given in Appendix B. In Section 3.2 we examine the action of the mapping on exceptional lines, with details given in Appendix C. We then describe approximate solutions near exceptional lines, with particular attention paid to a neighbourhood of the origin in Section 3.3. A conclusion and summary are provided in Section 4.

2. The Initial Value Space

In this section, we describe the major features of the initial value space $\mathcal{S}$, obtained by resolution of singularities of the dynamical system (1.2). The details of the calculations are provided in Appendices A and B.

2.1. Resolution of the Initial Value Space. Consider the dynamical system (1.2) in $\mathbb{P}^1 \times \mathbb{P}^1$, which is covered by four charts described in Figure 2.1. Singularities occur where the numerator and denominator vanish simultaneously in the definition of at least one of $\overline{u}$, $\overline{v}$, $\underline{u}$, or $\underline{v}$. We call these *base points*, following [**7**]. We resolve these in the standard way[2] to obtain successive charts (u_{ij}, v_{ij}), where i corresponds to the number of blow-ups and j corresponds to a chart indicated in Figure 2.1.

The definition of $\overline{v}$ and $\underline{u}$ in Equations (1.2) show that there are two base points in Chart 1:

$$b_1 : \begin{pmatrix} u_{01} \\ v_{01} \end{pmatrix} = \begin{pmatrix} t \\ 0 \end{pmatrix}, \qquad b_2 : \begin{pmatrix} u_{01} \\ v_{01} \end{pmatrix} = \begin{pmatrix} 0 \\ qt \end{pmatrix}, \tag{2.1}$$

while there is one base point in each of Charts 2 and 3 respectively:

$$b_3 : \begin{pmatrix} u_{02} \\ v_{02} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \qquad b_4 : \begin{pmatrix} u_{03} \\ v_{03} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{2.2}$$

The explicit resolutions of these base points are provided in Appendix A.

[2]We recall here that the standard operation of resolution or blowing up a base point at (α, β) means replacing (u, v) by two new coordinate charts $\tilde{u} = (u - \alpha)/(v - \beta), \tilde{v} = v - \beta$, or $\hat{u} = (u - \alpha), \hat{v} = (v - \beta)/(u - \alpha)$. This has the effect of replacing the base point by an *exceptional* line. Each resolution lowers the self-intersection number of the line containing the base point by unity. In the chart $(\tilde{u}, \tilde{v})$, this line is defined by $\tilde{v} = 0$ and parametrized by $\tilde{u}$, whilst in chart $(\hat{u}, \hat{v})$, the line is defined by $\hat{u} = 0$ and parameterized by $\hat{v}$. For this and other standard constructions of algebraic geometry, the reader is referred to [**11**].

Chart 3: $\begin{pmatrix} u_{03} \\ v_{03} \end{pmatrix} = \begin{pmatrix} u \\ 1/v \end{pmatrix}$ *Chart 4*: $\begin{pmatrix} u_{04} \\ v_{04} \end{pmatrix} = \begin{pmatrix} 1/u \\ 1/v \end{pmatrix}$



Chart 1: $\begin{pmatrix} u_{01} \\ v_{01} \end{pmatrix} = \begin{pmatrix} u \\ v \end{pmatrix}$ *Chart 2*: $\begin{pmatrix} u_{02} \\ v_{02} \end{pmatrix} = \begin{pmatrix} 1/u \\ v \end{pmatrix}$

FIGURE 2.1. The four coordinate charts of $\mathbb{P}^1 \times \mathbb{P}^1$

The resolution of b_1 and b_2 lead to no further singularities (see §A.1-A.2). However, the resolutions of b_3 and b_4 lead to four further base points, which are given in terms of coordinate charts defined in §A.3-A.4 by

$$b_5 : \begin{pmatrix} u_{31} \\ v_{31} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad b_6 : \begin{pmatrix} u_{42} \\ v_{42} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \tag{2.3}$$

$$b_7 : \begin{pmatrix} u_{51} \\ v_{51} \end{pmatrix} = \begin{pmatrix} -q \\ 0 \end{pmatrix}, b_8 : \begin{pmatrix} u_{61} \\ v_{61} \end{pmatrix} = \begin{pmatrix} -q \\ 0 \end{pmatrix}. \tag{2.4}$$

The resolution of b_3 leads to b_5, b_4 leads to b_6, while those of b_5 and b_6 lead to b_7 and b_8 respectively. The details can be found in §A.3-A.8. The exceptional lines E_j replacing each base point b_j, $j = 1, \ldots, 8$ are drawn schematically in Figure 2.2, where H_u denotes a line where u is constant and H_v denotes a line where v is constant.

2.2. Divisors. The equivalence classes of lines H_u, H_v, and E_i for $i = 1, .., 8$ form the standard basis of the Picard group $\mathrm{Pic}(\mathcal{S})$ [**11**]. This is equipped with a symmetric bilinear form $(\cdot, \cdot)$ on elements of $\mathrm{Pic}(\mathcal{S})$ given by

(1) $(H_u, H_u) = (H_v, H_v) = (H_u, E_i) = (H_v, E_i) = 0$,
(2) $(H_u, H_v) = 1$,
(3) $(E_i, E_j) = -\delta_{ij}$.



FIGURE 2.2. The rational surface $\mathcal{S}$ obtained from the resolution of singularities in Equation (1.2) indicating proper transforms of relevant curves.

From the resolution described above (see Figure 2.2), it follows that the nodal curves with self-intersection -2 are

$$\begin{aligned} D_1 &= H_v - E_1 - E_3, & D_5 &= H_v - E_4 - E_6, \\ D_2 &= E_3 - E_5, & D_6 &= E_6 - E_8, \\ D_3 &= E_5 - E_7, & D_7 &= E_4 - E_6, \\ D_4 &= H_u - E_3 - E_5, & D_8 &= H_u - E_2 - E_4. \end{aligned}$$

These form irreducible components of the anti-canonical divisor, defined by $-K_{\mathcal{S}} = 2H_u + 2H_v - \sum_{i=1}^{8} E_i$. By the definitions above, we have $-K_{\mathcal{S}} = \sum_{j=1}^{8} D_j$.

From the intersection rules on the Picard group given above, it follows that $(D_1, D_2) = 1 = (D_1, D_8)$, while $(D_1, D_j) = 0$ for $j = 3, 4, 5, 6, 7$. Further details about the intersections of D_j and the corresponding generalized Cartan matrix are provided in Appendix B. By representing each D_j as a node and connecting two nodes D_i, D_j by a simple edge if $(D_i, D_j) \neq 0$, we obtain the Dynkin diagram given in Figure 2.3, which represents the affine Weyl group $A_7^{(1)}$. By considering its orthogonal complement in $\mathrm{Pic}(\mathcal{S})$, we also show explicitly in Appendix B that Equation (1.2) has a symmetry group given by $A_1^{(1)}$.

3. Dynamics

Let the forward iteration given in Equation (1.2) on $\mathcal{S}$ be denoted by φ. We study iterations of points near the components D_j of the anti-canonical divisor in §3.1, near the exceptional lines E_1, E_2, E_7, E_8 in §3.2 and in a region around the origin punctured at E_1 and E_2 in §3.3. The section ends with the explicit linearisation of the map on the Picard lattice.



FIGURE 2.3. Intersection diagram of $\{D_j\}_{j=1}^8$, which is equivalent to the Dynkin diagram of $A_7^{(1)}$.

3.1. Dynamics near the irreducible components D_i. The iteration of these components under φ is computed in Appendix B. For example, consider $D_1 = H_u - E_1 - E_3$. Given $t = t_0$, a finite initial point near D_1 lies near the line $v_{01} = 0$ (with $u_{01} \neq t_0$), so that $v_{01}(t_0) = \delta v'_{01}(t_0) + O(\delta^2)$ for some finite $v'_{01}(t_0)$, and small $0 < \delta \ll 1$. The results of Appendix B show that

$$\begin{cases} u_{01}(t_0/q^8) & \sim & -\dfrac{u_{01}(t_0)\, t_0^3}{q^{14}}, \\ v_{01}(t_0/q^8) & \sim & \delta \dfrac{q v'_{01}(t_0) t_0^2 \left(q^6 + u_{01}(t_0) t_0^2\right)}{(u_{01}(t_0) - t_0)}, \end{cases} \tag{3.1}$$

which again lies near D_1 after iterating forward 4 steps. Arguments similar to these in Appendix B show that the set $D_1, \ldots, D_8$ is closed under the action of φ, the time evolution of D_j is periodic and, moreover, their iteration occurs in two orbits of period four each, as shown in Figure 3.1.



FIGURE 3.1. Action of φ on divisors.

3.2. Dynamics near exceptional lines E_1, E_2, E_7, E_8. As shown in Appendix C, we have the following behaviour under forward iteration

$$E_7 \to E_1 \to H_u + 2H_v - E_2 - E_3 - E_4 - E_5 - E_7, \quad \text{and} \quad E_2 \to E_8 \to H_v - E_2, \tag{3.2}$$

and under backward iteration we have

$$H_u - E_1 \leftarrow E_7 \leftarrow E_1, \quad \text{and} \quad 2H_u + H_v - E_1 - E_3 - E_4 - E_6 - E_8 \leftarrow E_2 \leftarrow E_8. \tag{3.3}$$

This means that under forward and backward iterations of these exceptional lines, we are sent eventually to regular points. There are certain special cases: for example $(u_{11}, v_{11}) = (0, 0)$ on E_1 is mapped under forward iteration to the point $(u_{71}, v_{71}) = (-q^2\xi, 0)$ on E_7. This point is mapped through the chain (3.2) once more, before emerging into regular space.

3.3. Dynamics near the origin. Here we consider dynamics in a neighbourhood of the origin in initial value space. Assume $d_{r_1}(0, 0)$ is a disk of radius r_1 around the origin in $\mathcal{S}$ for sufficiently small $r_1 > 0$. We assume that t lies in a small domain $d_{r_2}(t_0)$, around a given point $t_0 \ll 1$, for sufficiently small $r_2 > 0$ and work in the domain $U \subset d_{r_1}(0, 0)$ lying in $\mathcal{S}$, which is punctured at $(t, 0)$ and $(0, qt)$. We denote the small punctured domain around $(t, 0)$ by σ_1 and that around $(0, qt)$ by σ_2. An example of such a domain is drawn in Figure 3.2.



FIGURE 3.2. Punctured domain U

The mapping of σ_j, $j = 1, 2$ under φ are obtained by the results in Appendix C. These show that the exceptional lines in each region act as repellers, and that iterates move away from each region.

In the domain U, for sufficiently small (u, v), we find

$$\overline{u} \sim -\frac{u\,v}{t} \qquad\qquad \underline{u} \sim -\frac{qt}{uv^2} \tag{3.4}$$

$$\overline{v} \sim -\frac{t}{u^2 v} \qquad\qquad \underline{v} \sim -\frac{u\,v}{q\,t} \tag{3.5}$$

These show that U is mapped to a domain with large values of v (i.e., towards the origin in chart 3, which corresponds to $(u, v) = (0, \infty)$) by the forward map, while it is mapped to a domain with large values of u (i.e., towards the origin in chart 2, which corresponds to $(u, v) = (\infty, 0)$) by the backward map. In other words, under the forward map, a domain with large values of v and finite u is attracted to the origin, while values of u and v lying in U are repelled away to a domain with large values of u and finite v. That is, the origin is a saddle-point of the dynamical system. This result is consistent with the origin being a repelling fixed point of the autonomous map, and with the quicksilver solution being an unstable solution, as shown in [**12**].

3.4. Summary of behaviours. Appendix B contains explicit calculations of the mappings of the irreducible components D_j of the anti-canonical divisor, and Appendix C contains explicit calculations of the mappings of the remaining exceptional lines.

Based on these results, we can express the action of φ on elements of the standard basis of the Picard group in matrix form:

$$\varphi \begin{pmatrix} H_u \\ H_v \\ E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \\ E_6 \\ E_7 \\ E_8 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 0 & -1 & -1 & -1 & -1 & 0 & 0 & 0 \\ 2 & 3 & 0 & -2 & -2 & -1 & -1 & -1 & -1 & 0 \\ 1 & 2 & 0 & -1 & -1 & -1 & -1 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 0 & -1 & -1 & -1 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & -1 & -1 & -1 & -1 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & -1 & -1 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} H_u \\ H_v \\ E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \\ E_6 \\ E_7 \\ E_8 \end{pmatrix} \tag{3.6}$$

where $\varphi : \mathrm{Pic}(\mathcal{S}) \to \mathrm{Pic}(\mathcal{S})$ is the action of forward iteration in n. Because all the eigenvalues of this matrix are roots of unity, its *algebraic entropy* (see Bellon and Viallet [**3**]) vanishes. This shows that the complexity of the dynamics of this system does not grow exponentially in the time step n.

4. Conclusions

In this paper, we explicitly constructed the initial value space of the q-discrete first Painlevé system (1.2) as $t \to 0$, and for that of the scalar equation (1.1), in the limit $\xi \to \infty$. We provided the coordinate charts of the space of initial values in detail, identifying exceptional lines and irreducible components of the anti-canonical divisor and, based on these, showed how to deduce qualitative information about the dynamics in the limit.

We found that the union of exceptional lines is a repeller for the flow; i.e., if a solution is near an exceptional line at any given time, after one time step it is immediately mapped to a different region. This is analogous to the results for Painlevé I found in [**8**]. The set of the irreducible components of the anti-canonical divisor is invariant under the time flow and the dynamical system is periodic on the level of each such component.

Nevertheless, the solutions are neither simple or periodic. In particular, we showed that the origin acts as a saddle point for the generic flow. The remaining fixed point solutions, which approach the singularities on three fibres of the autonomous pencil (1.4) (see §1.1) are the discrete analogues of the so called *tronquée* solutions identified by Boutroux for the first Painlevé equation, which also approach double points of the latter's autonomous Hamiltonian.

The dynamics of most other q-discrete Painlevé equations in limits when $t \to 0$ or $t \to \infty$ remain unknown. In particular, it would be useful to study the q-discrete Painlevé equations known as $q\mathrm{P}_{\mathrm{II}}$ and $q\mathrm{P}_{\mathrm{III}}$, which have $q\mathrm{P}_{\mathrm{I}}$ as a degenerate limit.

Finally, since these q-discrete Painlevé equations arise as reductions of Yang-Baxter maps, it would be interesting to relate their limiting behaviours to the physical setting in which such maps play a role.

Appendix A. Blow up of base points

Here we provide explicit details of the process of resolution of each of the 8 base points of $q\mathrm{P_I}$.

A.1. Blow up of base point b_1. This base point arises in Chart 1 when we iterate the map forward.

Define new coordinates

$$\begin{pmatrix} u_{11} \\ v_{11} \end{pmatrix} = \begin{pmatrix} (u_{01}-t)/v_{01} \\ v_{01} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{01} \\ v_{01} \end{pmatrix} = \begin{pmatrix} u_{11}v_{11}+t \\ v_{11} \end{pmatrix}. \tag{A.1}$$

In these coordinates, Equations (1.2) become

$$\begin{cases} \overline{u} &= \dfrac{(u_{11}v_{11}+t)\left(qu_{11}-u_{11}^2v_{11}^2t-2u_{11}v_{11}t^2-t^3\right)}{qu_{11}^2} \\ \overline{v} &= \dfrac{u_{11}}{(u_{11}v_{11}+t)^{\,2}} \end{cases} \tag{A.2a}$$

and

$$\begin{cases} \underline{u} &= \dfrac{v_{11}-qt}{v_{11}^2\,(u_{11}v_{11}+t)} \\ \underline{v} &= \dfrac{v_{11}\,(u_{11}v_{11}+t)\left(v_{11}-q^2u_{11}v_{11}^3t-qt-q^2v_{11}^2t^2\right)}{(v_{11}-qt)^{\,2}} \end{cases} \tag{A.2b}$$

where the base point b_1 is replaced by the exceptional line E_1 defined by $v_{11}=0$. There are no new base points in this chart.

To look in the other chart, define new coordinates

$$\begin{pmatrix} u_{12} \\ v_{12} \end{pmatrix} = \begin{pmatrix} u_{01}-t \\ v_{01}/(u_{01}-t) \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{01} \\ v_{01} \end{pmatrix} = \begin{pmatrix} u_{12}+t \\ u_{12}v_{12} \end{pmatrix}. \tag{A.3}$$

The equations in these coordinates become

$$\begin{cases} \overline{u} &= \dfrac{(u_{12}+t)\,v_{12}\left(q-u_{12}^2v_{12}t-2u_{12}v_{12}t^2-v_{12}t^3\right)}{q} \\ \overline{v} &= \dfrac{1}{(u_{12}+t)^{\,2}v_{12}} \end{cases} \tag{A.4a}$$

and

$$\begin{cases} \underline{u} &= \dfrac{u_{12}v_{12}-qt}{u_{12}^2\,(u_{12}+t)\,v_{12}^2} \\ \underline{v} &= \dfrac{u_{12}\,(u_{12}+t)\,v_{12}\left(u_{12}v_{12}-q^2u_{12}^3v_{12}^2t-qt-q^2u_{12}^2v_{12}^2t^2\right)}{(u_{12}v_{12}-qt)^{\,2}} \end{cases}$$

where the base point b_1 is replaced by the exceptional line E_1 defined by $u_{12}=0$. There are no new base points in this chart.

A.2. Blow up of base point b_2. Define new coordinates

$$\begin{pmatrix} u_{21} \\ v_{21} \end{pmatrix} = \begin{pmatrix} u_{01}/(v_{01}-qt) \\ v_{01}-qt \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{01} \\ v_{01} \end{pmatrix} = \begin{pmatrix} u_{21}v_{21} \\ v_{21}+qt \end{pmatrix}. \tag{A.5}$$

The equations in these coordinates become

$$\left\{\begin{array}{rcl} \overline{u} &=& \dfrac{u_{21}v_{21}\left(v_{21}+qt\right)\left(qu_{21}v_{21}-qt-u_{21}^2v_{21}^3t-qu_{21}^2v_{21}^2t^2\right)}{q\left(u_{21}v_{21}-t\right)^2} \\ \overline{v} &=& \dfrac{u_{21}v_{21}-t}{u_{21}^2v_{21}^2\left(v_{21}+qt\right)} \end{array}\right. \tag{A.6a}$$

and

$$\left\{\begin{array}{rcl} \underline{u} &=& \dfrac{1}{u_{21}\left(v_{21}+qt\right)^2} \\ \underline{v} &=& u_{21}\left(v_{21}+qt\right)\left(1-q^2u_{21}v_{21}^2t-2q^3u_{21}v_{21}t^2-q^4u_{21}t^3\right) \end{array}\right. \tag{A.6b}$$

where the base point b_2 is replaced by the exceptional line E_2 defined by $v_{21}=0$. There are no new base points in this chart.

To look in the other chart, define new coordinates

$$\left(\begin{array}{c} u_{22} \\ v_{22} \end{array}\right)=\left(\begin{array}{c} u_{01} \\ (v_{01}-qt)/u_{01} \end{array}\right) \quad\Rightarrow\quad \left(\begin{array}{c} u_{01} \\ v_{01} \end{array}\right)=\left(\begin{array}{c} u_{22} \\ u_{22}v_{22}+qt \end{array}\right). \tag{A.7}$$

The equations in these coordinates become

$$\left\{\begin{array}{rcl} \overline{u} &=& \dfrac{u_{22}\left(u_{22}v_{22}+qt\right)\left(qu_{22}-qt-u_{22}^3v_{22}t-qu_{22}^2t^2\right)}{q\left(u_{22}-t\right)^2} \\ \overline{v} &=& \dfrac{u_{22}-t}{u_{22}^2\left(u_{22}v_{22}+qt\right)} \end{array}\right. \tag{A.8a}$$

and

$$\left\{\begin{array}{rcl} \underline{u} &=& \dfrac{v_{22}}{\left(u_{22}v_{22}+qt\right)^2} \\ \underline{v} &=& \dfrac{\left(u_{22}v_{22}+qt\right)\left(v_{22}-q^2u_{22}^2v_{22}^2t-2q^3u_{22}v_{22}t^2-q^4t^3\right)}{v_{22}^2} \end{array}\right. \tag{A.8b}$$

where E_2 is defined by $u_{22}=0$. There are no new base points in this chart.

A.3. Blow up of base point b_3. Define new coordinates

$$\left(\begin{array}{c} u_{31} \\ v_{31} \end{array}\right)=\left(\begin{array}{c} u_{02}/v_{02} \\ v_{02} \end{array}\right) \quad\Rightarrow\quad \left(\begin{array}{c} u_{02} \\ v_{02} \end{array}\right)=\left(\begin{array}{c} u_{31}v_{31} \\ v_{31} \end{array}\right). \tag{A.9}$$

In terms of the original variables, this is

$$\left(\begin{array}{c} u_{31} \\ v_{31} \end{array}\right)=\left(\begin{array}{c} 1/(uv) \\ v \end{array}\right) \quad\Rightarrow\quad \left(\begin{array}{c} u \\ v \end{array}\right)=\left(\begin{array}{c} 1/(u_{31}v_{31}) \\ v_{31} \end{array}\right). \tag{A.10}$$

The equations in these coordinates become

$$\left\{\begin{array}{rcl} \overline{u} &=& \dfrac{v_{31}\left(qu_{31}-qu_{31}^2v_{31}t-t\right)}{qu_{31}\left(1-u_{31}v_{31}t\right)^2} \\ \overline{v} &=& u_{31}\left(1-u_{31}v_{31}t\right) \end{array}\right. \tag{A.11a}$$

and

$$\left\{\begin{array}{rcl} \underline{u} &=& \dfrac{u_{31}\left(v_{31}-qt\right)}{v_{31}} \\ \underline{v} &=& \dfrac{u_{31}v_{31}-q^2v_{31}t-qu_{31}t}{u_{31}^2\left(v_{31}-qt\right)^2} \end{array}\right. \tag{A.11b}$$

where the base point b_3 is replaced by the exceptional line E_3 defined by $v_{31} = 0$. There is a new base point at

$$b_5 : \begin{pmatrix} u_{31} \\ v_{31} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{A.12}$$

To look in the other chart, define new coordinates

$$\begin{pmatrix} u_{32} \\ v_{32} \end{pmatrix} = \begin{pmatrix} u_{02} \\ v_{02}/u_{02} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{02} \\ v_{02} \end{pmatrix} = \begin{pmatrix} u_{32} \\ u_{32}v_{32} \end{pmatrix}. \tag{A.13}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{32} \\ v_{32} \end{pmatrix} = \begin{pmatrix} 1/u \\ uv \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 1/u_{32} \\ u_{32}v_{32} \end{pmatrix}. \tag{A.14}$$

The equations now become

$$\begin{cases} \overline{u} &= \dfrac{u_{32}v_{32}\,(q - qu_{32}t - v_{32}t)}{q\,(1 - u_{32}t)^{\,2}}, \\ \overline{v} &= \dfrac{1 - u_{32}t}{v_{32}}, \end{cases} \tag{A.15a}$$

and

$$\begin{cases} \underline{u} &= \dfrac{u_{32}v_{32} - qt}{u_{32}v_{32}^2}, \\ \underline{v} &= \dfrac{v_{32}\left(u_{32}v_{32} - q^2u_{32}v_{32}^2t - qt\right)}{(u_{32}v_{32} - qt)^{\,2}}. \end{cases} \tag{A.15b}$$

where E_3 is now given by $u_{32} = 0$. There are no further base points in this chart.

A.4. Blow up of base point b_4. Define new coordinates

$$\begin{pmatrix} u_{41} \\ v_{41} \end{pmatrix} = \begin{pmatrix} u_{03}/v_{03} \\ v_{03} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{03} \\ v_{03} \end{pmatrix} = \begin{pmatrix} u_{41}v_{41} \\ v_{41} \end{pmatrix}. \tag{A.16}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{41} \\ v_{41} \end{pmatrix} = \begin{pmatrix} uv \\ 1/v \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} u_{41}v_{41} \\ 1/v_{41} \end{pmatrix}. \tag{A.17}$$

The equations in these coordinates become

$$\begin{cases} \overline{u} &= \dfrac{u_{41}\left(qu_{41}v_{41} - qt - u_{41}^2v_{41}t\right)}{q\,(u_{41}v_{41} - t)^{\,2}} \\ \overline{v} &= \dfrac{u_{41}v_{41} - t}{u_{41}^2v_{41}} \end{cases} \tag{A.18a}$$

and

$$\begin{cases} \underline{u} &= \dfrac{1 - qv_{41}t}{u_{41}} \\ \underline{v} &= \dfrac{u_{41}v_{41}\left(1 - q^2u_{41}t - qv_{41}t\right)}{(1 - qv_{41}t)^{\,2}} \end{cases} \tag{A.18b}$$

where b_4 is replaced by the exceptional line E_4, which is defined by $v_{41} = 0$. There are no new base points in this chart.

To look in the other chart, define new coordinates

$$\begin{pmatrix} u_{42} \\ v_{42} \end{pmatrix} = \begin{pmatrix} u_{03} \\ v_{03}/u_{03} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{03} \\ v_{03} \end{pmatrix} = \begin{pmatrix} u_{42} \\ u_{42}v_{42} \end{pmatrix}. \tag{A.19}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{42} \\ v_{42} \end{pmatrix} = \begin{pmatrix} u \\ 1/(uv) \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} u_{42} \\ 1/(u_{42}v_{42}) \end{pmatrix}. \tag{A.20}$$

The equations now become

$$\begin{cases} \overline{u} &= \dfrac{qu_{42}v_{42} - qv_{42}t - u_{42}t}{qv_{42}^2\left(u_{42}-t\right)^2} \\ \overline{v} &= \dfrac{v_{42}\left(u_{42}-t\right)}{u_{42}} \end{cases} \tag{A.21a}$$

and

$$\begin{cases} \underline{u} &= v_{42}\left(1-qu_{42}v_{42}t\right) \\ \underline{v} &= \dfrac{u_{42}\left(v_{42}-q^2t-qu_{42}v_{42}^2t\right)}{v_{42}\left(1-qu_{42}v_{42}t\right)^2} \end{cases} \tag{A.21b}$$

where E_4 is defined by $u_{42} = 0$. There is a new base point in this chart at

$$b_6 : \begin{pmatrix} u_{42} \\ v_{42} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{A.22}$$

A.5. Blow up of base point b_5. Define new coordinates

$$\begin{pmatrix} u_{51} \\ v_{51} \end{pmatrix} = \begin{pmatrix} u_{31}/v_{31} \\ v_{31} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{31} \\ v_{31} \end{pmatrix} = \begin{pmatrix} u_{51}v_{51} \\ v_{51} \end{pmatrix}. \tag{A.23}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{51} \\ v_{51} \end{pmatrix} = \begin{pmatrix} 1/(uv^2) \\ v \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 1/(u_{51}v_{51}^2) \\ v_{51} \end{pmatrix}. \tag{A.24}$$

The equations in these coordinates become

$$\begin{cases} \overline{u} &= \dfrac{qu_{51}v_{51} - qu_{51}^2v_{51}^3t - t}{qu_{51}\left(1-u_{51}v_{51}^2t\right)^2} \\ \overline{v} &= u_{51}v_{51}\left(1-u_{51}v_{51}^2t\right) \end{cases} \tag{A.25a}$$

and

$$\begin{cases} \underline{u} &= u_{51}\left(v_{51}-qt\right) \\ \underline{v} &= \dfrac{u_{51}v_{51}-q^2t-qu_{51}t}{u_{51}^2v_{51}\left(v_{51}-qt\right)^2} \end{cases} \tag{A.25b}$$

where b_5 is replaced by the exceptional line E_5 defined by $v_{51} = 0$. There is a new base point at

$$b_7 : \begin{pmatrix} u_{51} \\ v_{51} \end{pmatrix} = \begin{pmatrix} -q \\ 0 \end{pmatrix}. \tag{A.26}$$

To look in the other chart, define new coordinates

$$\begin{pmatrix} u_{52} \\ v_{52} \end{pmatrix} = \begin{pmatrix} u_{31} \\ v_{31}/u_{31} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{31} \\ v_{31} \end{pmatrix} = \begin{pmatrix} u_{52} \\ u_{52}v_{52} \end{pmatrix}. \tag{A.27}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{52} \\ v_{52} \end{pmatrix} = \begin{pmatrix} 1/(uv) \\ uv^2 \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} 1/(u_{52}^2v_{52}) \\ u_{52}v_{52} \end{pmatrix}. \tag{A.28}$$

The equations in these coordinates become

$$\begin{cases} \overline{u} &= \dfrac{v_{52}\left(qu_{52}-qu_{52}^3v_{52}t-t\right)}{q\left(1-u_{52}^2v_{52}t\right)^2} \\ \overline{v} &= u_{52}\left(1-u_{52}^2v_{52}t\right) \end{cases} \tag{A.29a}$$

and

$$\begin{cases} \underline{u} &= \dfrac{u_{52}v_{52}-qt}{v_{52}} \\ \underline{v} &= \dfrac{u_{52}v_{52}-q^2v_{52}t-qt}{u_{52}\left(u_{52}v_{52}-qt\right)^2} \end{cases} \tag{A.29b}$$

whereas E_5 is now $u_{52}=0$. There are no further base points appearing here.

A.6. Blow up of base point b_6. Define new coordinates

$$\begin{pmatrix} u_{61} \\ v_{61} \end{pmatrix} = \begin{pmatrix} u_{42}/v_{42} \\ v_{42} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{42} \\ v_{42} \end{pmatrix} = \begin{pmatrix} u_{61}v_{61} \\ v_{61} \end{pmatrix}. \tag{A.30}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{61} \\ v_{61} \end{pmatrix} = \begin{pmatrix} u^2v \\ 1/(uv) \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} u_{61}v_{61} \\ 1/(u_{61}v_{61}^2) \end{pmatrix}. \tag{A.31}$$

The equations in these new coordinates become

$$\begin{cases} \overline{u} &= \dfrac{qu_{61}v_{61}-qt-u_{61}t}{qv_{61}\left(u_{61}v_{61}-t\right)^2} \\ \overline{v} &= \dfrac{u_{61}v_{61}-t}{u_{61}} \end{cases} \tag{A.32a}$$

and

$$\begin{cases} \underline{u} &= v_{61}\left(1-qu_{61}v_{61}^2t\right) \\ \underline{v} &= \dfrac{u_{61}\left(v_{61}-q^2t-qu_{61}v_{61}^3t\right)}{\left(1-qu_{61}v_{61}^2t\right)^2} \end{cases} \tag{A.32b}$$

where the base point b_6 is replaced by the exceptional line E_6 defined by $v_{61}=0$. There is a new base point at

$$b_8 : \begin{pmatrix} u_{61} \\ v_{61} \end{pmatrix} = \begin{pmatrix} -q \\ 0 \end{pmatrix}. \tag{A.33}$$

To look in the other chart, define new coordinates

$$\begin{pmatrix} u_{62} \\ v_{62} \end{pmatrix} = \begin{pmatrix} u_{42} \\ v_{42}/u_{42} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{42} \\ v_{42} \end{pmatrix} = \begin{pmatrix} u_{62} \\ u_{62}v_{62} \end{pmatrix}. \tag{A.34}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{62} \\ v_{62} \end{pmatrix} = \begin{pmatrix} u \\ 1/(u^2v) \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} u_{62} \\ 1/(u_{62}^2v_{62}) \end{pmatrix}. \tag{A.35}$$

The equations in these new coordinates become

$$\begin{cases} \overline{u} &= \dfrac{qu_{62}v_{62}-qv_{62}t-t}{qu_{62}v_{62}^2\left(u_{62}-t\right)^2} \\ \overline{v} &= v_{62}\left(u_{62}-t\right) \end{cases} \tag{A.36a}$$

and

$$\left\{\begin{array}{lcl} \underline{u} &=& u_{62}v_{62}\left(1-qu_{62}^2v_{62}t\right) \\ \underline{v} &=& \dfrac{\left(u_{62}v_{62}-q^2t-qu_{62}^3v_{62}^2t\right)}{v_{62}\left(1-qu_{62}^2v_{62}t\right)^2} \end{array}\right. \tag{A.36b}$$

where E_6 is now defined by $u_{62}=0$. There are no further base points in this chart.

A.7. Blow up of base point b_7. Define new coordinates

$$\left(\begin{array}{c} u_{71} \\ v_{71} \end{array}\right)=\left(\begin{array}{c} (u_{51}+q)/v_{51} \\ v_{51} \end{array}\right) \quad\Rightarrow\quad \left(\begin{array}{c} u_{51} \\ v_{51} \end{array}\right)=\left(\begin{array}{c} u_{71}v_{71}-q \\ v_{71} \end{array}\right). \tag{A.37}$$

In terms of the original variables, this is

$$\left(\begin{array}{c} u_{71} \\ v_{71} \end{array}\right)=\left(\begin{array}{c} (1+quv^2)/(uv^3) \\ v \end{array}\right) \quad\Rightarrow\quad \left(\begin{array}{c} u \\ v \end{array}\right)=\left(\begin{array}{c} 1/[(u_{71}v_{71}-q)v_{71}^2] \\ v_{71} \end{array}\right). \tag{A.38}$$

The equations now become

$$\left\{\begin{array}{lcl} \overline{u} &=& \dfrac{q^2v_{71}-qu_{71}v_{71}^2+q^3v_{71}^3t-2q^2u_{71}v_{71}^4t+qu_{71}^2v_{71}^5t+t}{q\left(q-u_{71}v_{71}\right)\left(1+qv_{71}^2t-u_{71}v_{71}^3t\right)^2} \\ \overline{v} &=& v_{71}\left(u_{71}v_{71}-q\right)\left(1+qv_{71}^2t-u_{71}v_{71}^3t\right), \end{array}\right. \tag{A.39}$$

and

$$\left\{\begin{array}{lcl} \underline{u} &=& -\left(q-u_{71}v_{71}\right)\left(v_{71}-qt\right) \\ \underline{v} &=& \dfrac{-q+u_{71}v_{71}-qu_{71}t}{\left(u_{71}v_{71}-q\right)^2\left(v_{71}-qt\right)^2} \end{array}\right. \tag{A.40}$$

where the base point b_7 is now replaced by the exceptional line E_7 defined by $v_{71}=0$. There are no further base points in this chart.

To look in the other chart, define new coordinates

$$\left(\begin{array}{c} u_{72} \\ v_{72} \end{array}\right)=\left(\begin{array}{c} u_{51}+q \\ v_{51}/(u_{51}+q) \end{array}\right) \quad\Rightarrow\quad \left(\begin{array}{c} u_{51} \\ v_{51} \end{array}\right)=\left(\begin{array}{c} u_{72}-q \\ u_{72}v_{72} \end{array}\right). \tag{A.41}$$

In terms of the original variables, this is

$$\left(\begin{array}{c} u_{72} \\ v_{72} \end{array}\right)=\left(\begin{array}{c} (1+quv^2)/(uv^2) \\ uv^3/(1+quv^2) \end{array}\right) \quad\Rightarrow\quad \left(\begin{array}{c} u \\ v \end{array}\right)=\left(\begin{array}{c} 1/[(u_{72}-q)u_{72}^2v_{72}^2] \\ u_{72}v_{72} \end{array}\right). \tag{A.42}$$

The equations in these coordinates become

$$\left\{\begin{array}{lcl} \overline{u} &=& \dfrac{q^2u_{72}v_{72}-qu_{72}^2v_{72}+q^3u_{72}^3v_{72}^3t-2q^2u_{72}^4v_{72}^3t+qu_{72}^5v_{72}^3t+t}{q\left(q-u_{72}\right)\left(1+qu_{72}^2v_{72}^2t-u_{72}^3v_{72}^2t\right)^2} \\ \overline{v} &=& u_{72}v_{72}\left(u_{72}-q\right)\left(1+qu_{72}^2v_{72}^2t-u_{72}^3v_{72}^2t\right) \end{array}\right. \tag{A.43}$$

and

$$\left\{\begin{array}{lcl} \underline{u} &=& -\left(q-u_{72}\right)\left(u_{72}v_{72}-qt\right) \\ \underline{v} &=& \dfrac{-qv_{72}+u_{72}v_{72}-qt}{v_{72}\left(u_{72}-q\right)^2\left(u_{72}v_{72}-qt\right)^2} \end{array}\right. \tag{A.44}$$

where now E_7 is defined by $u_{72}=0$. There are no further base points appearing here.

A.8. Blow up of base point b_8. Define new coordinates

$$\begin{pmatrix} u_{81} \\ v_{81} \end{pmatrix} = \begin{pmatrix} (u_{61}+q)/v_{61} \\ v_{61} \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{61} \\ v_{61} \end{pmatrix} = \begin{pmatrix} u_{81}v_{81}-q \\ v_{81} \end{pmatrix}. \tag{A.45}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{81} \\ v_{81} \end{pmatrix} = \begin{pmatrix} (u^2v+q)uv \\ 1/(uv) \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} (u_{81}v_{81}-q)v_{81} \\ 1/[(u_{81}v_{81}-q)v_{81}^2] \end{pmatrix}. \tag{A.46}$$

The equations become

$$\begin{cases} \overline{u} &= \dfrac{qu_{81}v_{81}-q^2-u_{81}t}{q\left(qv_{81}-u_{81}v_{81}^2+t\right)^2} \\ \overline{v} &= \dfrac{qv_{81}-u_{81}v_{81}^2+t}{q-u_{81}v_{81}t} \end{cases} \tag{A.47}$$

and

$$\begin{cases} \underline{u} &= v_{81}\left(1+q^2v_{81}^2t-qu_{81}v_{81}^3t\right) \\ \underline{v} &= \dfrac{(u_{81}v_{81}-q)\left(v_{81}+q^2v_{81}^3t-q^2t-qu_{81}v_{81}^4t\right)}{\left(1+q^2v_{81}^2t-qu_{81}v_{81}^3t\right)^2} \end{cases} \tag{A.48}$$

where the base point b_8 is now replaced by the exceptional line E_8 defined by $v_{81}=0$. There are no further base points in this chart.

To look in the other chart, define new coordinates

$$\begin{pmatrix} u_{82} \\ v_{82} \end{pmatrix} = \begin{pmatrix} u_{61}+q \\ v_{61}/(u_{61}+q) \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u_{61} \\ v_{61} \end{pmatrix} = \begin{pmatrix} u_{82}-q \\ u_{82}v_{82} \end{pmatrix}. \tag{A.49}$$

In terms of the original variables, this is

$$\begin{pmatrix} u_{82} \\ v_{82} \end{pmatrix} = \begin{pmatrix} u^2v+q \\ 1/[(u^2v+q)uv] \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} (u_{82}-q)u_{82}v_{82} \\ 1/[(u_{82}-q)u_{82}^2v_{82}^2] \end{pmatrix}. \tag{A.50}$$

The equations in these coordinates become

$$\begin{cases} \overline{u} &= \dfrac{qu_{82}v_{82}-q^2v_{82}-t}{qv_{82}\left(qu_{82}v_{82}-u_{82}^2v_{82}+t\right)^2} \\ \overline{v} &= \dfrac{qu_{82}v_{82}-u_{82}^2v_{82}+t}{q-u_{82}} \end{cases} \tag{A.51}$$

and

$$\begin{cases} \underline{u} &= u_{82}v_{82}\left(1+q^2u_{82}^2v_{82}^2t-qu_{82}^3v_{82}^2t\right) \\ \underline{v} &= \dfrac{(u_{82}-q)\left(u_{82}v_{82}+q^2u_{82}^3v_{82}^3t-q^2t-qu_{82}^4v_{82}^3t\right)}{\left(1+q^2u_{82}^2v_{82}^2t-qu_{82}^3v_{82}^2t\right)^2} \end{cases} \tag{A.52}$$

where E_8 is now defined by $u_{82}=0$. There are no further base points in this chart.

Appendix B. Dynamics of solutions near divisors D_i

Defining $A_{ij} := 2(D_i,D_j)/(D_j,D_j)$, we can express the intersection information between D_i and D_j in a generalised Cartan matrix $A := (A_{ij})_{i,j=1}^8$. For

$\{D_j\}_{j=1}^8$ defined in §2.2, we have

$$A = \begin{pmatrix} 2 & -1 & 0 & 0 & 0 & 0 & 0 & -1 \\ -1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 2 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 2 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ -1 & 0 & 0 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}.$$

B.1. Symmetries of the system. We find the symmetries of the system by constructing vectors orthogonal to the components $D_1, \ldots, D_8$, and defining corresponding actions which leave this set invariant. A vector $\alpha \in \mathrm{Pic}(\mathcal{S})$ is given by the linear combination

$$\alpha := \alpha_u H_u + \alpha_v H_v + \sum_{i=1}^{8} \alpha_i E_i \tag{B.1}$$

and its intersection with each D_j is given by

$$\begin{aligned} (\alpha, D_1) &= \alpha_u + \alpha_1 + \alpha_3, \\ (\alpha, D_2) &= -\alpha_3 + \alpha_5, \\ (\alpha, D_3) &= -\alpha_5 + \alpha_7, \\ (\alpha, D_4) &= \alpha_v + \alpha_3 + \alpha_5, \\ (\alpha, D_5) &= \alpha_u + \alpha_4 + \alpha_6, \\ (\alpha, D_6) &= -\alpha_6 + \alpha_8, \\ (\alpha, D_7) &= -\alpha_4 + \alpha_6, \\ (\alpha, D_8) &= \alpha_v + \alpha_2 + \alpha_4. \end{aligned}$$

For orthogonality to be satisfied, it follows that

$$\alpha_3 = \alpha_5 = \alpha_7 \quad =: \quad a, \tag{B.2}$$

$$\alpha_4 = \alpha_6 = \alpha_8 \quad =: \quad b, \tag{B.3}$$

where a and b are arbitrary, and hence

$$\alpha_u = -2b, \tag{B.4}$$

$$\alpha_v = -2a, \tag{B.5}$$

$$\alpha_1 = 2b - a, \tag{B.6}$$

$$\alpha_2 = 2a - b. \tag{B.7}$$

Thus α becomes

$$\begin{aligned} \alpha &= a(-2H_v - E_1 + 2E_2 + E_3 + E_5 + E_7) + b(-2H_u + 2E_1 - E_2 + E_4 + E_6 + E_8) \\ &=: aF_1 + bF_2 \end{aligned} \tag{B.8}$$

where, if we define $B_{i,j} := 2(F_i, F_j)/(F_j, F_j)$, we find the generalised Cartan matrix $B := (B_{ij})_{i,j=1}^2$ given by

$$B = \begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}. \tag{B.9}$$

This leads to the Dynkin diagram shown in Figure B.1, which corresponds to the root lattice $A_1^{(1)}$.



FIGURE B.1. Dynkin diagram corresponding to the symmetry group.

For each vector F_i, we define an action on an element $x \in \mathrm{Pic}(X)$ as follows:

$$\begin{aligned} w_{F_i}(x) &:= x - 2\frac{(x,F_i)}{(F_i,F_i)}F_i \\ &= x + \frac{1}{4}(x,F_i)F_i. \end{aligned} \tag{B.10}$$

The action of w_{F_1} on $\mathrm{Pic}(X)$ is

$$w_{F_1}\begin{pmatrix} H_u \\ H_v \\ E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \\ E_6 \\ E_7 \\ E_8 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1/2 & -1 & -1/2 & 0 & -1/2 & 0 & -1/2 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1/2 & 3/4 & 1/2 & 1/4 & 0 & 1/4 & 0 & 1/4 & 0 \\ 0 & 1 & 1/2 & 0 & -1/2 & 0 & -1/2 & 0 & -1/2 & 0 \\ 0 & 1/2 & 1/4 & -1/2 & 3/4 & 0 & -1/4 & 0 & -1/4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1/2 & 1/4 & -1/2 & -1/4 & 0 & 3/4 & 0 & -1/4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1/2 & 1/4 & -1/2 & -1/4 & 0 & -1/4 & 0 & 3/4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} H_u \\ H_v \\ E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \\ E_6 \\ E_7 \\ E_8 \end{pmatrix}, \tag{B.11}$$

and the action of w_{F_2} on $\mathrm{Pic}(X)$ is

$$w_{F_2}\begin{pmatrix} H_u \\ H_v \\ E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \\ E_6 \\ E_7 \\ E_8 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & -1 & 1/2 & 0 & -1/2 & 0 & -1/2 & 0 & -1/2 \\ 1 & 0 & 0 & 1/2 & 0 & -1/2 & 0 & -1/2 & 0 & -1/2 \\ -1/2 & 0 & 1/2 & 3/4 & 0 & 1/4 & 0 & 1/4 & 0 & 1/4 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1/2 & 0 & -1/2 & 1/4 & 0 & 3/4 & 0 & -1/4 & 0 & -1/4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1/2 & 0 & -1/2 & 1/4 & 0 & -1/4 & 0 & 3/4 & 0 & -1/4 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 1/2 & 0 & -1/2 & 1/4 & 0 & -1/4 & 0 & -1/4 & 0 & 3/4 \end{pmatrix} \begin{pmatrix} H_u \\ H_v \\ E_1 \\ E_2 \\ E_3 \\ E_4 \\ E_5 \\ E_6 \\ E_7 \\ E_8 \end{pmatrix}. \tag{B.12}$$

Note that the action of w_{F_i} on each of the D_i is the identity, i.e.,

$$w_{F_i}(D_j) = D_j \quad \forall i = 1,2; \quad j = 1,..,8, \tag{B.13}$$

and taking either of these actions to the second power gives the identity, so the w_{F_i} are indeed reflections; the span of these reflections forms the affine Weyl group W. Including a Dynkin automorphism $\sigma : F_1 \leftrightarrow F_2$, we obtain the extended affine Weyl group $\widetilde{W}$. Note that the action $(w_{F_2}w_{F_1})^4$ has the same effect on the elements of the Picard group as does the 4th power of the mapping (1.1). That is, if we denote the operation of forward shift in the discrete time variable by φ, we have

$$\varphi^4(K) = (w_{F_2}w_{F_1})^4 K, \tag{B.14}$$

for K an element of the Picard group.

In the following, we examine behaviour of solutions near each irreducible component $D_1, \ldots, D_8$ of the anti-canonical divisor. We focus on the forward iteration here, as the case of backward iteration is entirely analogous.

B.2. Behaviour near $D_1 := H_{v=0} - E_1 - E_3$. The component D_1 is essentially the coordinate axis $v_{01} = v_{02} = 0$, where $u_{01} \neq t, u_{02} \neq 0$. Suppose that at a time t_1 we are near D_1, i.e., $v_{01}(t_1) = v_{02}(t_1)$ is close to zero. Expanding the equations (1.2) for forward iteration in the chart (u_{41}, v_{41}), we find to leading order for $v_{01}(t_1) = \epsilon \ll 1$

$$\left\{ \begin{array}{rcl} \overline{u}_{41}(t_1) & = & \dfrac{1}{u_{01}(t_1)} + \mathcal{O}(\epsilon) \\ \overline{v}_{41}(t_1) & \sim & \dfrac{u_{01}(t_1)^2 v_{01}(t_1)}{u_{01}(t_1) - t_1} = \mathcal{O}(\epsilon) \end{array} \right. \tag{B.15}$$

while for $v_{02}(t_1) = \epsilon$ we have

$$\left\{ \begin{array}{rcl} \overline{u}_{41}(t_1) & = & u_{02}(t_1) + \mathcal{O}(\epsilon) \\ \overline{v}_{41}(t_1) & \sim & \dfrac{v_{02}(t_1)}{u_{02}(t_1)^2(1 - u_{02}(t_1)t_1)} = \mathcal{O}(\epsilon). \end{array} \right. \tag{B.16}$$

The image lies near the line $v_{41} = 0$, that is, near the component D_7.

B.3. Behaviour near D_2. The component D_2 is defined in local coordinates by $v_{31} = 0$, $u_{32} = 0$, where $u_{31} \neq 0$. Suppose that at a time t_2 we are near this divisor; i.e., $v_{31}(t_2), u_{32}(t_2)$ are close to zero. Expanding the equations (1.2) for forward iteration in the chart (u_{31}, v_{31}), we find to leading order for $v_{31}(t_2) = \epsilon \ll 1$

$$\left\{ \begin{array}{rcl} \overline{u}_{01}(t_2) & \sim & \dfrac{v_{31}(t_2)(q u_{31}(t_2) - t_2)}{q u_{31}(t_2)} = \mathcal{O}(\varepsilon) \\ \overline{v}_{01}(t_2) & = & u_{31}(t_2) + \mathcal{O}(\varepsilon) \end{array} \right. \tag{B.17}$$

and for $u_{32}(t_2) = \varepsilon$

$$\left\{ \begin{array}{rcl} \overline{u}_{01}(t_2) & \sim & \dfrac{v_{32}(t_2) u_{32}(t_2)(q - v_{32}(t_2)t_2)}{q} = \mathcal{O}(\varepsilon) \\ \overline{v}_{01}(t_2) & = & \dfrac{1}{v_{32}(t_2)} + \mathcal{O}(\varepsilon). \end{array} \right. \tag{B.18}$$

We see that the image near the line $u_{01} = 0$, near the component D_8.

B.4. Behaviour near D_3. The component D_3 is the line $E_5 - E_7$ is defined in local coordinates by $v_{51} = 0, u_{52} = 0$, where $u_{51} \neq -q$. Suppose that at a time t_3 we are near this component; i.e., $v_{51}(t_3) = \varepsilon$, or $u_{52}(t_3) = \varepsilon$ where $\varepsilon \ll 1$. Expanding the equations (1.2), we see that in the coordinates of the first chart

$$\left\{ \begin{array}{rcl} \overline{u}_{01}(t_3) & = & -\dfrac{1}{q t_3 u_{51}(t_3)} + \mathcal{O}(\varepsilon), \\ \overline{v}_{01}(t_3) & \sim & u_{51}(t_3) v_{51}(t_3) = \mathcal{O}(\varepsilon), \end{array} \right. \tag{B.19}$$

while, in the coordinates of the second chart we obtain

$$\left\{ \begin{array}{rcl} \overline{u}_{01}(t_3) & = & -\dfrac{v_{52}(t_3)}{q t_3} + \mathcal{O}(\varepsilon), \\ \overline{v}_{01}(t_3) & \sim & u_{52}(t_3) = \mathcal{O}(\varepsilon). \end{array} \right. \tag{B.20}$$

The image lies near the line $v_{01} = 0$, or rather the component D_1.

B.5. Behaviour near D_4. The component D_4 is given by $u_{02} = u_{04} = 0$, where $v_{02} \neq 0$. Suppose that at a time t_4 we are near this component; i.e., $u_{02}(t_4) = u_{04}(t_4) = \varepsilon \ll 1$. Expanding the equations for u_{31}, v_{31}, we find in the first chart

$$\begin{cases} \overline{u}_{31}(t_4) &= -\dfrac{qt_4}{v_{02}(t_4)} + \mathcal{O}(\varepsilon) \\ \overline{v}_{31}(t_4) &\sim \dfrac{u_{02}(t_4)}{v_{02}(t_4)} = \mathcal{O}(\varepsilon) \end{cases} \tag{B.21}$$

while in the coordinates of the second chart we obtain

$$\begin{cases} \overline{u}_{31}(t_4) &= -qt_4v_{04}(t_4) + \mathcal{O}(\varepsilon) \\ \overline{v}_{31}(t_4) &\sim v_{04}(t_4)u_{04}(t_4) = \mathcal{O}(\varepsilon). \end{cases} \tag{B.22}$$

The image lies near the line $v_{31} = 0$, or rather the component D_2.

B.6. Behaviour near D_5. The divisor D_5 is essentially the coordinate axis $v_{03} = v_{04} = 0$, where $u_{03} \neq 0$. Suppose that at a time t_5 we are near this divisor; i.e., $v_{03}(t_5) = v_{04}(t_5) = \varepsilon$ is close to zero. Expanding the equations for u_{51}, v_{51}, we find for $|v_{03}| \ll 1$

$$\begin{cases} \overline{u}_{51}(t_5) &= -qt_5u_{03}(t_5) + \mathcal{O}(\varepsilon) \\ \overline{v}_{51}(t_5) &\sim \dfrac{v_{03}(t_5)(u_{03}(t_5) - t_5)}{u_{03}(t_5)^2} = \mathcal{O}(\varepsilon) \end{cases} \tag{B.23}$$

while for $|v_{04} \ll 1$, we find

$$\begin{cases} \overline{u}_{51}(t_5) &= -\dfrac{qt_5}{u_{04}(t_5)} + \mathcal{O}(\varepsilon) \\ \overline{v}_{51}(t_5) &\sim u_{04}(t_5)v_{04}(t_5)(1 - u_{04}(t_5)t_5) + \mathcal{O}(\varepsilon) \end{cases} \tag{B.24}$$

The image lies near the line $v_{51} = 0$, or rather the component D_3

B.7. Behaviour near D_6. The component D_6 is defined in local coordinates by $v_{61} = 0, u_{62} = 0$, where $u_{61} \neq -q$. Suppose that at a time t_6 we are near this component; i.e., $v_{61}(t_6)$, $u_{62}(t_6)$ are close to zero. Expanding the equations for forward iteration in chart 2, we see that for $|v_{61}| = \varepsilon \ll 1$

$$\begin{cases} \overline{u}_{02}(t_6) &\sim -\dfrac{qv_{61}(t_6)}{t_6(q + u_{61}(t_6))} = \mathcal{O}(\varepsilon), \\ \overline{v}_{02}(t_6) &= -\dfrac{1}{t_6u_{61}(t_6)} + \mathcal{O}(\varepsilon), \end{cases} \tag{B.25}$$

and for $|u_{62}| = \varepsilon \ll 1$,

$$\begin{cases} \overline{u}_{02}(t_6) &\sim -\dfrac{qv_{62}(t_6)^2u_{62}(t_6)}{t_6(qv_{62}(t_6) + 1)} = \mathcal{O}(\varepsilon), \\ \overline{v}_{02}(t_6) &= -\dfrac{v_{62}(t_6)}{t_6} + \mathcal{O}(\varepsilon). \end{cases} \tag{B.26}$$

We see that this is near the line $u_{02} = 0$, or rather the component D_4.

B.8. Behaviour near D_7. The component D_7 is the line $E_4 - E_6$ that arises under the first blow-up at the point b_4 in the region where u is finite, v is infinite. It is defined in local coordinates by $v_{41} = 0$, $u_{42} = 0$, where $v_{42} \neq 0$. Suppose that at a time t_7 we are near this component; i.e., $v_{41}(t_7) = \varepsilon$, $u_{42}(t_7) = \varepsilon$, where $\varepsilon \ll 1$. Expanding the equations in chart 3, we see that in the coordinates of the first chart

$$\begin{cases} \overline{u}_{03}(t_7) & \sim & -t_7 u_{41}(t_7) + \mathcal{O}(\varepsilon) \\ \overline{v}_{03}(t_7) & = & -t_7 u_{41}(t_7)^2 v_{41}(t_7) = \mathcal{O}(\varepsilon) \end{cases} \tag{B.27}$$

and in the coordinates of the second chart

$$\begin{cases} \overline{u}_{03}(t_7) & = & -\dfrac{t_7}{v_{42}(t_7)} + \mathcal{O}(\varepsilon), \\ \overline{v}_{03}(t_7) & \sim & -\dfrac{t_7 u_{42}(t_7)}{v_{42}(t_7)} = \mathcal{O}(\varepsilon). \end{cases} \tag{B.28}$$

The image lies near the line $v_{03} = 0$, or rather the component D_5.

B.9. Behaviour near D_8. The component D_8 is essentially the coordinate axis $u_{01} = u_{03} = 0$ where $v_{01} \neq qt, v_{03} \neq 0$. Suppose that at a time t_8 we are near this component; i.e., $u_{01}(t_8) = u_{03}(t_8) = \varepsilon \ll 1$. Expanding the equations for forward iteration in (u_{61}, v_{61}), we see that in the coordinates of the first chart

$$\begin{cases} \overline{u}_{61}(t_8) & = & -t_8 v_{01}(t_8) + \mathcal{O}(\varepsilon), \\ \overline{v}_{61}(t_8) & \sim & u_{01}(t_8) = \mathcal{O}(\varepsilon) \end{cases} \tag{B.29}$$

and in the coordinates of the second chart

$$\begin{cases} \overline{u}_{61}(t_8) & = & -\dfrac{t_8}{v_{03}(t_8)} + \mathcal{O}(\varepsilon) \\ \overline{v}_{61}(t_8) & \sim & u_{03}(t_8) = \mathcal{O}(\varepsilon). \end{cases} \tag{B.30}$$

The image lies near the line $v_{61} = 0$, or rather the component D_6.

Appendix C. Mappings of remaining exceptional lines

In this appendix, we analyse the dynamics starting with initial values near the exceptional lines E_1, E_2, E_7, E_8. For simplicity and conciseness, we focus on the local results of iterating a neighbourhood near each exceptional line and, to do so, we assume that initial values in a neighbourhood of E_i are analytic functions of t close to a point t_i, for each $i = 1, 2, 7, 8$. Being analytic away from singularities, the birational map (1.2) maps a disk near an exceptional line to another disk of non-zero size.

C.1. Behaviour near E_1. The exceptional line E_1 arises under blow-up of the point b_1 in the region where u, v are finite; it is the line $v_{11} = 0$, or equivalently, $u_{12} = 0$. Expanding the equations (1.2) for forward and backward iteration, we find in the coordinates of the first chart:

$$\begin{cases} \overline{u}_{01}(t_1) & \sim & t_1 \dfrac{q u_{11}(t_1) - t_1^3}{q u_{11}(t_1)^2}, \\ \overline{v}_{01}(t_1) & \sim & \dfrac{u_{11}(t_1)}{t_1^2}, \end{cases} \qquad \begin{cases} \underline{u}_{71}(t_1) & \sim & -qt_1(1 + q u_{11}(t_1)), \\ \underline{v}_{71}(t_1) & \sim & -\dfrac{v_{11}(t_1)}{q}, \end{cases} \tag{C.1}$$

and in the coordinates of the second chart:

$$\begin{cases} \overline{u}_{01}(t_1) &\sim \dfrac{t_1 v_{12}(t_1)(q - v_{12}(t_1)t_1^3)}{q}, \\ \overline{v}_{01}(t_1) &\sim \dfrac{1}{t_1^2 v_{12}(t_1)}, \end{cases} \qquad \begin{cases} \underline{u}_{71}(t_1) &\sim -\dfrac{q(q + v_{12}(t_1))}{t_1 v_{12}(t_1)}, \\ \underline{v}_{71}(t_1) &\sim -\dfrac{v_{12}(t_1)u_{12}(t_1)}{q}. \end{cases} \tag{C.2}$$

The image under the forward map is a curve in regular space. Under backward iteration, we are mapped to near the line $v_{71} = 0$, i.e., to E_7.

C.2. Behaviour near E_2. The exceptional line E_2 is the line $v_{21} = 0$, $u_{22} = 0$. Suppose that at a time t_2 we are near this exceptional line; i.e., $v_{21}(t_2)$, $u_{22}(t_2)$ are close to zero. Expanding the equations for (u_{81}, v_{81}) expressed in terms of the coordinates (u_{21}, v_{21}), we find

$$\begin{cases} \overline{u}_{81}(t_2) &\sim -\dfrac{(qu_{21}(t_2) + 1)}{t_2 u_{21}(t_2)}, \\ \overline{v}_{81}(t_2) &\sim u_{21}(t_2)v_{21}(t_2), \end{cases} \qquad \begin{cases} \underline{u}_{01}(t_2) &\sim \dfrac{t_2^2}{q^2 u_{21}(t_2)}, \\ \underline{v}_{01}(t_2) &\sim qt_2 u_{21}(t_2)(1 - q^4 t_2^3 u_{21}(t_2), \end{cases} \tag{C.3}$$

while in the coordinates of the second chart, we obtain

$$\begin{cases} \overline{u}_{81}(t_2) &\sim -\dfrac{q + v_{22}(t_2)}{t_2}, \\ \overline{v}_{81}(t_2) &\sim u_{22}(t_2), \end{cases} \qquad \begin{cases} \underline{u}_{01}(t_2) &\sim \dfrac{v_{22}(t_2)}{q^2 t_2^2}, \\ \underline{v}_{01}(t_2) &\sim \dfrac{qt_2(v_{22}(t_2) - q^4 t_2^3)}{v_{22}(t_2)^2}. \end{cases} \tag{C.4}$$

We see that under forward iteration the image lies near the line $v_{81} = 0$, i.e., near E_8. Under backward iteration, this is a curve in regular space.

C.3. Behaviour near E_7. E_7 is the line $v_{71} = 0$ or equivalently $u_{72} = 0$. Suppose that at a time t_7 we are near this exceptional line; i.e., $v_{71}(t_7)$, $u_{72}(t_7)$ are close to zero. Expanding the equations for (u_{11}, v_{11}) in coordinates (u_{71}, v_{71}), we find:

$$\begin{cases} \overline{u}_{11}(t_7) &\sim -\dfrac{q^3 + u_{71}(t_7)t_7}{q^4}, \\ \overline{v}_{11}(t_7) &\sim -qv_{71}(t_7), \end{cases} \tag{C.5}$$

while the backward map for (u, v) in (u_{71}, v_{71}) gives

$$\begin{cases} \underline{u}_{01}(t_7) &\sim q^2 t_7, \\ \underline{v}_{01}(t_7) &\sim -\dfrac{(1 + t_7 u_{71}(t_7))}{q^3 t_7^2}, \end{cases} \tag{C.6}$$

Now consider these maps in in coordinates (u_{72}, v_{72}):

$$\begin{cases} \overline{u}_{11}(t_7) &\sim -\dfrac{q^3 v_{72}(t_7) + t_7}{q^4 v_{72}(t_7)}, \\ \overline{v}_{11}(t_7) &\sim -qv_{72}(t_7)u_{72}(t_7), \end{cases} \tag{C.7}$$

$$\begin{cases} \underline{u}_{01}(t_7) &\sim q^2 t_7, \\ \underline{v}_{01}(t_7) &\sim -\dfrac{(t_7 + v_{72}(t_7))}{q^3 t_7^2 v_{72}(t_7)}. \end{cases} \tag{C.8}$$

Under forward iteration the image lies near the line $v_{11} = 0$, or rather E_1 shifted forward. Under backward iteration, we are mapped to near the line $u_{01} = q^2 t$.

C.4. Behaviour near E_8. The exceptional line E_8 is the line $v_{81} = 0$, $u_{82} = 0$. Suppose that at a time t_8, we are near this exceptional line; i.e., $v_{81}(t_8)$, $u_{82}(t_8)$ are close to zero. Expanding the forward and backward maps in chart 1 for small $v_{81}(t_8)$, $u_{82}(t_8)$, we find in coordinates (u_{81}, v_{81})

$$\begin{cases} \overline{u}_{01}(t_8) & \sim & -\dfrac{(q^2 + t_8\, u_{81}(t_8))}{qt_8^2}, \\ \overline{v}_{01}(t_8) & \sim & \dfrac{t_8}{q}, \end{cases} \qquad \begin{cases} \underline{u}_{21}(t_8) & \sim & -\dfrac{1}{q\,(1 + qt_8u_{81})}, \\ \underline{v}_{21}(t_8) & \sim & -\,q\,(1 + qt_8u_{81})v_{81}, \end{cases} \tag{C.9}$$

while in the coordinates of the second chart

$$\begin{cases} \overline{u}_{01}(t_8) & \sim & -\dfrac{q^2v_{82}(t_8) + 1)}{qt_8^2v_{82}(t_8)}, \\ \overline{v}_{01}(t_8) & \sim & \dfrac{t_8}{q}, \end{cases} \qquad \begin{cases} \underline{u}_{21}(t_8) & \sim & -\dfrac{v_{82}(t_8)}{q(q\,t_8 + v_{82})}, \\ \underline{v}_{21}(t_8) & \sim & q\,(q\,t_8 + v_{82})u_{82}. \end{cases} \tag{C.10}$$

We see that under forward iteration this is near the line $v_{01} = t/q$. Under backward iteration, we are mapped to near the line $v_{21} = 0$, or rather E_2 shifted backward.

Acknowledgements

The authors would like to thank H. Dullin, A. Dzhamay, C. Lustri and T. Takenawa for informative discussions. The research reported in this paper was supported by Australian Laureate Fellowship Grant #FL120100094 from the Australian Research Council.

References

1. V. E. Adler, A. Bobenko, and Y. B. Suris. *Classification of integrable equations on quad-graphs. The consistency approach.* Comm. Math. Phys. **233** (2003) 513–543.
2. J.A. Atkinson and Y. Yamada, *Quadrirational Yang-Baxter maps and the affine-E_8 Painlevé lattice*, arXiv:1405.2745 [nlin.SI] (2014).
3. M. P. Bellon and C.-M. Viallet, *Algebraic entropy*, Comm. Math. Phys. **204** (1999) 425–437
4. C. M. Bender, S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers.* McGraw-Hill, New York, 1978.
5. P. Boutroux, *Recherches sur les transcendantes de M. Painlevé et l'étude asymptotique des équations différentielles du second ordre*, Ann. École Norm. **30** (1913) 265–375.
6. B. Dubrovin, T. Grava, C. Klein, *On universality of critical behavior in the focusing nonlinear Schrödinger equation, elliptic umbilic catastrophe and the Tritronquée Solution to the Painlevé equation.* J. Nonlin. Sci **19** (2009) 57–94.
7. J.J. Duistermaat, *Discrete integrable systems: QRT maps and elliptic surfaces*, Springer, New York, 2010.
8. J.J. Duistermaat and N. Joshi, *Okamoto's space for the first Painlevé equation in Boutroux coordinates*, Arch. Rat. Mechs and Anal. **202** (2011) 707–785.
9. A.S. Fokas, A.R. Its and A.V. Kitaev, *Discrete Painlevé equations and their appearance in quantum gravity*, Comm. Math. Phys. **142** (1991) 313–344.
10. D. J. Gross and A. Migdal, *Nonperturbative two-dimensional quantum gravity*, Phys. Rev. Letts **64** (2) (1990) 127–130.
11. Robin Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer, 1977.
12. N. Joshi, *Quicksilver solutions of a q-difference first Painlevé equation*, Stud. Appl. Math **134** (2014) 233 – 251.
13. N. Joshi and A.V. Kitaev *On Boutrouxs tritronqueé solutions of the first Painlevé equation* Stud. Appl. Math. **107** (2001) 253.
14. N. Joshi, N. Nakazono, and Y. Shi *Geometric reductions of ABS equations on an n-cube to discrete Painlevé systems.* J. Phys. A **47** (2014) 505201 (16pp).

15. S. Nishioka, *Transcendence of solutions of q-Painlevé equation of type* $A_7^{(1)}$, Aequationes Mathematicae **79** (2010), no. 1-2, 1–12 (English).
16. Y. Ohyama, *Expansions on special solutions of the first q-Painlevé equation around the infinity*, Proceedings of the Japan Academy, Series A, Mathematical Sciences **86** (2010), no. 5, 91–92.
17. K. Okamoto, *Sur les feuilletages associés aux équation du second ordre à points critiques fixes de P. Painlevé. Espaces de conditions initiales.* Japanese J. Math. **5** (1979) 1–79.
18. V.G. Papageorgiou, A.G. Tongas, and A.P. Veselov. *Yang-Baxter maps and symmetries of integrable equations on quad-graphs.* J. Math. Phys. **47** (2006) 083502 (16pp).
19. H. Sakai, *Rational surfaces associated with affine root systems and geometry of the Painlevé equations*, Comm. Math. Phys. **220** (2001), no. 1, 165–229.
20. T. Tsuda, *Integrable mappings via rational elliptic surfaces*, J. Phys. A: Math. Gen. **37** (2004) 2721–2730

School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia
E-mail address: nalini.joshi@sydney.edu.au

School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia
E-mail address: sarah.lobb@sydney.edu.au